

## ABOUT THE NEOTROPICAL GENUS *LOPHEUCOILA* WELD (HYMENOPTERA: CYNIPOIDEA: FIGITIDAE): IDENTITY AND GEOGRAPHICAL DISTRIBUTION<sup>1</sup>

Fabiana E. Gallardo,<sup>2</sup> Norma B. Díaz,<sup>2</sup> and Jorge A. Guimarães<sup>3</sup>

**ABSTRACT:** In this paper a new species synonymy is established, *Lopheucoila anastrephae* (Rohwer, 1919) (= *Lopheucoila truncicola* Weld, 1951) and a new record of *Lopheucoila mexicana* Weld, 1951 is reported from Cuba.

**KEY WORDS:** *Zaeucoila* group, *Lopheucoila*, new record, new synonymy, parasitoids, frugivorous flies.

Díaz and Gallardo (1997, 1998a, b) and Gallardo and Díaz (1999) proposed the *Zaeucoila* group to be placed in the following neotropical genera: *Agrostocynips* Díaz, *Dettmeria* Borgmeier, *Moneucoela* Kieffer, *Penteucoila* Weld, *Rhabdeucoela* Kieffer, *Tropideucoila* Ashmead, *Zaeucoila* Ashmead and *Lopheucoila* Weld. Buffington and Scheffer (2008) have added four genera to the *Zaeucoila* group: *Dicerataspis* Ashmead, *Aegeseucoela* Buffington, *Moritiella* Buffington and *Preseucoela* Buffington. The *Zaeucoila* group species are koinobiont endoparasitoids of cyclorhaphous Diptera, parasitizing either leafminers (Agromyzidae) or the frugivorous flies of Drosophilidae, Lonchaeidae, Otitidae and Tephritidae (Borgmeier, 1935; Buffington, 2002, 2004; De Santis et al., 1976; Díaz and Valladares, 1979; Díaz et al., 1999; Fontal and Nieves-Aldrey, 2004; Guimarães and Zucchi, 2004; Murphy and La Salle, 1999; Rohwer, 1919; Wharton et al., 1998). Some of the frugivorous species are known as true “fruit flies” (Tephritidae) and are considered a most important group of pests of economic importance due to their damage to worldwide pomiculture (Aluja, 1994).

*Lopheucoila* is a Neotropical genus included in the *Zaeucoila* group. This genus was created by Weld (1951) based on *Diglyphosema anastrephae* Rohwer, 1919 (type species). More recently another two species, *L. mexicana* Weld and *L. truncicola* Weld, were also included in *Lopheucoila*. Thus, today, *Lopheucoila* is known to be represented by three species, which are associated with Lonchaeidae and Tephritidae larvae (Guimarães et al., 2003). However, Gallardo and Díaz (1999) reviewed this genus and observed that the differences between *L. truncicola* and *L. anastrephae*, mainly related to the body size and to the form of the edge of the posterior margin of the cup, were considered to be individual variations. This statement suggested that *L. truncicola* could be a synonym of *L. anastrephae*.

<sup>1</sup> Received on August 27, 2008. Accepted on December 21, 2008.

<sup>2</sup> División Entomología, Museo de La Plata, Paseo del Bosque, 1900 La Plata, Argentina. E-mail (FEG): gallardo@museo.fcnym.unlp.edu.ar, (NBD) ndiaz@fcnym.unlp.edu.ar

<sup>3</sup> EMBRAPA, Centro Nacional de Pesquisa de Hortalizas, Rodovia Brasília/Anápolis, Brazil 060, Km 09 CEP 70351-970, Gama – DF. E-mail: jorge@cnpat.embrapa.br

The objectives of this paper are to provide diagnostic characters for identification of the known species, to establish the appropriate nomenclatural changes, and update information on geographical distributions.

## METHODS

We studied 288 (144 females and 144 males) specimens belonging to *Lopheucoila*. The names of the institutions where the studied specimens are housed were designated by their initials between parentheses. The material belongs to the following institutions: Texas A & M University (TAMU), USA; United States National Museum of Natural History Washington (USNM), USA; Museu de Entomologia da Escola Superior de Agricultura "Luiz de Queiróz" (ESALQ), BRAZIL; Instituto Superior de Entomología de la Fundación Miguel Lillo (INSUE); ARGENTINA and Museo de La Plata (MLP), ARGENTINA.

All specimens were examined using a Leica S8APO stereomicroscope. Drawing was made with camera lucida adapted to a Leitz compound microscope.

The geographical distribution of the species following the scheme proposed by Cabrera and Willink (1980).

### *Lopheucoila* Weld (Figs. 1, 2)

*Lopheucoila* Weld, 1951: 223; 1952: 105, 195 (citation); Nordlander, 1982: 22 (citation); Díaz and Gallardo, 1997: 32 (citation); 1998a: 113 (citation), 1998b: 318 (citation); Wharton et al., 1998: 110 (distribution and bionomics); Gallardo and Díaz, 1999: 16 (review); Guimarães et al., 1999: 263 (bionomics), 2000: 129, 130, 133 (citation), 2003: 4 (distribution); Ovruski et al., 2000: 84 (citation); Uchôa-Fernandes et al., 2003: 182 (citation); Fontal-Cazalla et al., 2002: 186 (phylogeny); Fontal and Nieves-Aldrey, 2004: 64, 74 (citation); Strikis and do Prado, 2004: 647 (bionomics); García and Corseuil, 2004: 517, 520 (bionomics).

**Type Species.** *Diglyphosema anastrephae* Rohwer, 1919, by original designation of Weld, 1951.

**Distribution** (Fig. 1). Mexico, Costa Rica, Panama, Venezuela, Trinidad and Tobago, Peru, Brazil and Argentina (Gallardo and Díaz, 1999). Here we present a new record from Cuba. The geographical distribution of *Lopheucoila* species is exclusively Neotropical, mainly in biogeographic provinces characterized by wide variety of flora, such as the Xerofila Mexicana, Caribe (Domain Caribe), Amazonica, Pacifica, Cerrado, Paraense, Sabana, Atlántica (Domain Amazonico) and Altoandina (Domain Andino-Patagonico) *sensu* Cabrera and Willink (1980).

**Biology.** Koinobiont endoparasitoids of the third instar larvae-pupae of the dipterous families Lonchaeidae and Tephritidae.




Figure 1. Distribution of the species of *Lopheucoila*. Symbols indicate the localities.

*Lopheucoila mexicana* Weld, 1951 (Fig. 1)

*Lopheucoila mexicana* Weld, 1951: 224; Gallardo and Díaz, 1999: 21 (redescription).

**Diagnosis.** Keel of mesoscutum conspicuous. Base of metasoma smooth.

**Distribution.** Neotropical region (Fig. 1): Mexico (Gallardo and Díaz, 1999) and Cuba (**new record**).




Figure 2. Dorsal habitus of *Lopheucoila anastrephae* (Rohwer, 1919). Scale: 1mm.

**Hosts.** Unknown.

**Type Material.** MEXICO. Cordoba. Holotype Female, without date, Crawford col., (USNM Type 60981).

**Other material examined.** CUBA. Santiago, Pr. Gran Piedra 1100 m. Meteo Station, 1 female, 6-7-XII-1995; 1 female, 14-17-XII-1995, coll. L. Masner, (yellow pan traps) (TAMU).

*Lopheucoila anastrephae* (Rohwer, 1919) (Figs. 1, 2)

*Diglyphosema anastrephae* Rohwer, 1919: 156.

*Lopheucoila anastrephae*; Weld, 1951: 224; Wharton et al., 1998: 110 (distribution and bionomics); Gallardo and Díaz, 1999: 21 (redescription); Guimarães et al., 1999: 263 (bionomics), 2003: 4 (distribution); Ovruski et al., 2000: 84 (citation); Fontal-Cazalla et al., 2002: 186 (phylogeny); Uchôa-Fernandes et al., 2003: 182 (citation); Fontal and Nieves-Aldrey, 2004: 64, 74 (citation); Strikis and do Prado, 2004: 647 (bionomics); García and Corseuil, 2004: 517, 520 (bionomics).

*Lopheucoila truncicola* Weld, 1951: 223; Gallardo and Díaz, 1999: 23 (redescription); **New Synonymy.**

**Diagnosis.** Keel of mesoscutum inconspicuous on the distal part. Base of metasoma with two striate lateral depressions.

**Distribution.** Neotropical region (Fig. 1): Mexico, Costa Rica, Panama, Venezuela, Trinidad and Tobago, Brazil, Peru and Argentina (Gallardo and Díaz, 1999).

**Hosts.** Diptera-Tephritidae: *Anastrepha* sp. (Rohwer, 1919), *A. fraterculus* (Wied.) (Weld, 1951), *A. pseudoparallela* (Loew) and *A. amita* Zucchi (Guimarães et al., 2000). Diptera-Lonchaeidae: *Neosilba batesi* (Curran) (Wharton et al., 1998), *Neosilba* spp. (Guimarães et al., 1999, 2000; García and Corseuil, 2004), *Neosilba glaberrima* (Wied.) and *N. zadolicha* McAlpine and Steyskal (Strikis and do Prado, 2004).

**Type Material.** **TRINIDAD and TOBAGO.** Holotype Female of *L. anastrephae*, VII-1917, coll. Urich, (USNM Type 22029). **BRAZIL. Rio de Janeiro. Nova Iguassu,** Holotype Female of *L. truncicola*, 16-XI-1931, colls. Kisliuk and Cooley, (USNM Type 60980).

**Other material examined.** **ARGENTINA. Tucumán.** 1 female, 01-III-1991, obtained from puparium of Tephritidae in guava; **Horco Molle,** 3 females and 1 male, 28-III-1993, obtained from *Anastrepha* sp. in guava, coll. Ovruski, (INSUE). **BRAZIL. Mato Grosso do Sul. Anastácio, Sítio Carandazal,** 2 males, 30-III-1993; 1 male, 6-IV-1993; 1 female, 27-IV-1993; 1 female and 1 male, 18-V-1993; 6 females and 3 males, 24-V-1993; 1 female and 1 male, 1-VI-1993; 1 male, 21-VI-1993; 1 male, 25-VII-1993; 2 males, 2-VIII-1993; 2 males, 17-VIII-1993; 1 female and 4 males, 23-VIII-1993; 2 males, 30-VIII-1993; 1 female and 2 males, 6-IX-1993; 10 females and 1 male, 13-IX-1993; 2 females and 1 male, 27-IX-1993; 1 female, 4-X-1993; 10 females and 4 males, 1-XI-1993; 1 male, 25-XI-1993; 2 females and 1 male, 6-XII-1993; 1 male, 4-IV-1994; 1 female, 18-IV-1994; 1 male, 2-V-1994; 3 males, 16-V-1994; 1 male, 6-VI-1994; 1 male, 22-VIII-1994; 1 male, 9-IV-1995; 1 female, 16-IV-1995; 3 females and 3 males, 21-V-1995 (ESALQ); 3 females and 5 males, 28-V-1995 (MLP); **Sítio Santo Antônio,** 2 males, 25-IV-1993; **Colônia Jamic, Terenos,** 3 females and 17 males, 31-III-1993, 3 females and 2 males, 26-IV-1993; 2 males, 3-V-1993; 7 females and 6 males, 14-V-1993; 2 females and 2 males, 7-VI-1993; 3 females and 8 males, 22-III-1994; 1 female, 17-V-1994; 1 female, 12-VII-1994; 1 female, 14-I-1994; **Aquidauana, Fazenda Arardry,** 13 females and 2 males, 24-VI-1994; **Ranchinho Rochedo,** 1 female, 24-I-1994, coll. Uchoa F.; **Minas Gerais, Grupiara,** 1 males, 8-X-2000, coll. Guimarães; **São Paulo. Regente Feijó,** 3 males, 27-XI-1996; **São Bento do Sapucaí,** 14 females and 11 males, 1-III-1996; **Campinas,** 5 males, 3-II-1995; 2 males, 25-VI-1997; 18 females and 8 males, 20-XI-1997; 2 females and 2 males, 11-II-1998; 1 male, 9-V-1997, 3 males, 18-I-1996; 5 males, 30-VII-1997. **Assis,** 3 males, 5-II-1998; **Garça,** 1 male, 13-VIII-1997; **Limeira,** 1 female, 20-IV-1998, coll. M.F. Souza Filho (ESALQ); **Monte Alegre do sul. Estação experimental do Instituto Agrônômico,** 3 females and 1 male, 21-V-2002, 1 male, 23-IX-2002, 1 female, 8-X-2002, 1 female, 5-XI-2002, obtained from puparium of *Neosilba zadolicha* in "níspero," 3 females, 3-IX-2002, obtained from puparium of *Neosilba zadolichain* in peach, 1 male, 3-IV-2003, obtained from puparium of *Neosilba zadoli-*

*cha* in guava, coll. Strikis (MLP). **COSTA RICA. San José. San Ignacio de Acosta**, 10 females and 5 males, 20-II-1981, orange, coll. M. Fischel; 3 females and 3 males, 1979-1981, Gilstrap Medfly Project (TAMU).

**Remarks.** The study of a great number of specimens from a wide geographical distribution, allowed to us corroborate the hypothesis proposed by Gallardo and Díaz (1999), which established that the morphological differences between *Lopheucoila anastrephae* and *L. truncicola* are intraspecific variations and are not strong enough for them to be considered as two different species.

### ACKNOWLEDGEMENTS

The authors thank D. Smith and D. Furth (NMNH) for loaning the type material, E. Riley (TAMU), R.A. Zucchi (ESALQ) and M. V. Colomo de Correa (INSUE) for loaning us material for study. Cecilia Gorretta for technical support. This study was supported by Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and Universidad Nacional de La Plata (UNLP) – Argentina.

### LITERATURE CITED

- Aluja, M.** 1994. Bionomics and management of *Anastrepha*. Annual Review of Entomology 39: 155- 178.
- Borgmeier, T.** 1935. Sobre alguns Cynipidos parasiticos e cecidogenos do Brasil (Hymenoptera: Cynipidae). Archivos do Instituto de Biologia Vegetal 2(1): 97-124.
- Buffington, M. L.** 2002. Description of *Aegeseucoela* Buffington, new name, with notes on the status of *Gronotoma* Foerster (Hymenoptera: Figitidae: Eucoilinae). Proceedings of the Entomological Society of Washington 104(3): 589-601.
- Buffington, M. L.** 2004. The description of *Preseucoela* Buffington, new genus, with notes on the status of Nearctic species of *Agrostocynips* Díaz (Hymenoptera: Figitidae: Eucoilinae). Zootaxa 408: 1-11.
- Buffington, M. L. and S. J. Scheffer.** 2008. North American species of *Agrostocynips* Díaz (Hymenoptera: Figitidae: Eucoilinae), parasitoids of Agromyzidae (Diptera): bionomics and taxonomy. Zootaxa 1817: 39-48.
- Cabrera, A. and A. Willink.** 1980. Biogeografía de América Latina. Monografía Número. 13 Serie Biología. Organización de Estados Americanos. Washington, D.C. U.S.A. 122 pp.
- De Santis, L., N. B. Díaz, and I. Redolfi.** 1976. La mosca del girasol (Diptera, Agromyzidae) y sus himenópteros parasitoides. Dusenía 9(1): 31-38.
- Díaz, N. B. and G. Valladares.** 1979. Nota sobre *Agrostocynips clavatus* y los agromicidos hospedantes (Hymenoptera Cynipoidea). Neotropica 25(73): 23-26.
- Díaz, N. B. and F. E. Gallardo.** 1997. Revisión sistemática de las especies del género *Zaeucoila* (Hymenoptera, Cynipoidea, Eucoilidae). Revista Nicaragüense de Entomología 39: 31- 40.
- Díaz, N. B. and F. E. Gallardo.** 1998a. Revisión sistemática del género *Moneucoela* (Hymenoptera: Figitidae). Revista de la Sociedad Entomológica Argentina 57(1-4): 113-115.
- Díaz, N. B. and F. E. Gallardo.** 1998b. Revisión sistemática del género *Lopheucoila* Weld (Hymenoptera, Figitidae). p. 31. In, IV Congreso Argentino de Entomología. Mar del Plata, Argentina. 113 pp.
- Díaz, N. B., F. E. Gallardo, R. Monteiro, and M. Barbosa da Silva.** 1999. Himenópteros cinipoides parasitoides de dípteros agromicidos en Argentina y Brasil. p. 205. In, X Jornadas Fitosanitarias Argentinas. Jujuy, Argentina. 298 pp.

- Fontal, F. and J. L. Nieves-Aldrey.** 2004. Estudio comparado de diversidad de eucoilinos paleárticos (El Ventorrillo, España) y neotropicales (P. N. Coiba, Panamá) (Hymenoptera, Cynipoidea, Figitidae, Eucoilinae). *Boletín de la Sociedad Entomológica Aragonesa* 35: 51-101.
- Fontal-Cazalla, F. M., M. L. Buffington, G. Nordlander, J. Liljeblad, P. Ros-Farré, J. L. Nieves-Aldrey, J. Pujade-Villar, and F. Ronquist.** 2002. Phylogeny of the Eucoilinae (Hymenoptera: Cynipoidea: Figitidae). *Cladistics* 18: 154-199.
- Gallardo, F. E. and N. B. Díaz.** 1999. Revisión sistemática de las especies del género *Lopheucoila* Weld (Figitidae: Eucoilinae). *Revista Nicaragüense de Entomología* 47: 15-23.
- García, F. R. M. and E. Corseuil.** 2004. Native hymenopteran parasitoids associated with fruit flies (Diptera: Tephritidae) in Santa Catarina State, Brazil. *Florida Entomologist* 87: 517-521.
- Guimarães, J. and R. Zucchi.** 2004. Parasitism behavior of three species of Eucoilinae (Hymenoptera: Cynipoidea: Figitidae) fruit fly parasitoids (Diptera) in Brazil. *Neotropical Entomology* 33(2): 217-224.
- Guimarães, J., N. Díaz, and R. Zucchi.** 2000. Parasitóides (Figitidae: Eucoilinae). Chapter 16. pp. 127-134. In, A. Malavasi and R. A. Zucchi (Editor). *Moscas- das- frutas de importância econômica no Brasil, Conhecimento básico e aplicado*. Holos Editora. Ribeirão Preto, São Paulo, Brasil. 327 pp.
- Guimarães, J. A., F. E. Gallardo, N. B. Díaz, and R. A. Zucchi.** 2003. Eucoilinae species (Hymenoptera: Cynipoidea: Figitidae) parasitoids of fruit-infesting dipterous larvae in Brazil: identity, geographical distribution and host associations. *Zootaxa* 278: 1-23.
- Guimarães, J., R. Zucchi, N. Díaz, M. Souza Filho, and M. Uchôa F.** 1999. Espécies de Eucoilinae (Hymenoptera: Cynipoidea: Figitidae) parasitóides de larvas frugívoras (Diptera: Tephritidae e Lonchaeidae) no Brasil. *Anais da Sociedade Entomológica do Brasil* 28(2): 263-273.
- Murphy, S. T. and J. La Salle.** 1999. Balancing biological control strategies in the IPM of New World invasive *Liriomyza* leafminers in field vegetable crops. *Biocontrol News and Information* 20(3): 91-104.
- Nordlander, G.** 1982. Systematics and phylogeny of an interrelated group of genera within the family Eucoilidae (Insecta, Hymenoptera, Cynipoidea). Ph.D. Thesis Stockholm University, Zoology Department (Stockholm, Sweden). 34 pp.
- Ovruski, S., M. Aluja, J. Sivinski, and R. Wharton.** 2000. Hymenopteran parasitoids on fruit-infesting Tephritidae (Diptera) in Latin America and the southern United States: diversity, distribution, taxonomic status and their use in fruit fly biological control. *Integrated Pest Management Reviews* 5: 81-107.
- Rohwer, S.** 1919. Description of a new cynipid from Trinidad. *Proceedings of the Entomological Society of Washington* 21: 156.
- Strikis, P. and A. do Prado.** 2004. Parasitóides de *Neosilba zadolicha* e *N. glaberrima* (Diptera: Lonchaeidae Acalyptatae) em goiaba *Psidium guajaba* e jiló *Solanum gilo*. p. 647. In, *Resumos XX Congresso Brasileiro de Entomologia*. Gramado, Brasil. 695 pp.
- Uchôa- Fernandes, M. A., R. M. da S. Molina, I. De Oliveira, R. A. Zucchi, N. A. Canal, and N. B. Díaz.** 2003. Larval endoparasitoids (Hymenoptera) of frugivorous flies (Diptera, Tephritoidea) reared from fruits of the cerrado of the State of Mato Grosso do Sul, Brazil. *Revista Brasileira de Entomologia* 47(2): 181-186.
- Weld, L.** 1951. New Eucoilinae (Hymenoptera: Cynipoidea). *Proceedings of the Entomological Society of Washington* 53(4): 223-224.
- Weld, L.** 1952. *Cynipoidea (Hymenoptera) 1905-1950*. Ann Arbor, Michigan, U.S.A. 351 pp. (Privately printed.)
- Wharton, R., S. Ovruski, and F. Gilstrap.** 1998. Neotropical Eucoilidae (Cynipoidea) associated with fruit-infesting Tephritidae, with new records from Argentina, Bolivia and Costa Rica. *Journal of Hymenoptera Research* 7(1): 102-115.